

Brage IMR – *Havforskningsinstituttets institusjonelle arkiv*

Dette er forfatters siste versjon av den fagfelleverderte artikkelen, vanligvis omtalt som postprint. I Brage IMR er denne artikkelen ikke publisert med forlagets layout fordi forlaget ikke tillater dette. Du finner lenke til forlagets versjon i Brage-posten. Det anbefales at referanser til artikkelen hentes fra forlagets side.

Ved lenking til artikkelen skal det lenkes til post i Brage IMR, ikke direkte til pdf-fil.

Brage IMR – *Institutional repository of the Institute of Marine Research*

This is the author's last version of the article after peer review and is not the publisher's version, usually referred to as postprint. You will find a link to the publisher's version in Brage IMR. It is recommended that you obtain the references from the publisher's site.

Linking to the article should be to the Brage-record, not directly to the pdf-file.



**Effects of different modified diets on growth, digestive enzyme activities
and muscle compositions in juvenile Siamese fighting fish
(*Betta splendens* Regan, 1910)**

Karun Thongprajukaew^{a, b, #}, Uthaiwan Kovitvadhi^{b, c, *}, Satit Kovitvadhi^d,
Pisamai Somsueb^e, Krisna Rungruangsak-Torrissen^{b, f, **}

^a *Inter Departmental Multidisciplinary Graduate Program in Bioscience, Faculty of Science,
Kasetsart University, Bangkok 10900, Thailand*

^b *Biochemical Research Unit for Feed Utilization Assessment, Faculty of Science,
Kasetsart University, Bangkok 10900, Thailand*

^c *Department of Zoology, Faculty of Science, Kasetsart University, Bangkok 10900, Thailand*

^d *Department of Agriculture, Faculty of Science and Technology, Bansomdejchaopraya
Rajabhat University, Bangkok 10600, Thailand*

^e *Inland Fisheries Research and Development Bureau, Department of Fisheries,
Bangkok 10900, Thailand*

^f *Institute of Marine Research, Ecosystem Processes Research Group,
Matre Research Station, N-5984 Matredal, Norway*

* Corresponding author: Tel.: +66 2562 5444x3250; fax: +66 2562 5444x3202.

E-mail address: fsciutk@ku.ac.th (U. Kovitvadhi)

** Corresponding author: Tel.: +47 56367539; fax: +47 56367585.

E-mail address: Krisnart@imr.no (K. Rungruangsak-Torrissen)

Present address: Department of Applied Science, Faculty of Science

Prince of Songkla University, Songkhla, Thailand

Abstract

The effects of four modified diets (gamma-irradiated, microwave-irradiated, probiotic-supplemented and carbohydrases-supplemented diets) were studied on digestive enzyme specific activities and growth performance quality of juvenile Siamese fighting fish (*Betta splendens* Regan, 1910) during 2 weeks of critical and intensive rearing period. The modified procedures did not change biochemical compositions and gross energy of diets, but generally resulted in relatively higher *in vitro* digestibilities of protein and carbohydrate and fish survival rate, albeit insignificant. Only gamma irradiation significantly increased *in vitro* protein digestibility of the diet, and microwave irradiation increased starch gelatinization and water solubility ($P < 0.05$). Fish fed microwave-treated diet showed highest values in all studied growth indicators and digestive enzyme specific activities (except lipase), with significantly higher amylase specific activity and activity ratio of amylase to trypsin (A/T ratio). Correlation analysis indicated significant relationships ($P < 0.05$) among the levels of total protease, amylase and trypsin, and between SGR and A/T ratio. Muscle and body compositions of juveniles fed on microwave- or gamma- irradiated diets were similar to the control, while the juveniles fed on probiotic- or carbohydrases-supplemented diets showed lower protein depositions ($P < 0.05$). Similar levels of RNA, RNA/Protein ratio, and Protein/Lipid ratio in body and muscle in all dietary groups fed *ad libitum* suggested that the improved growth performance in juvenile Siamese fighting fish fed on microwave-irradiated diet may not be only due to improved physicochemical properties of the diet but also improved fish consumption rate.

Keywords: Digestive enzymes; *In vitro* digestibility; Modified diet; Muscle composition; Nutrient utilization; Siamese fighting fish

1. Introduction

Production of Siamese fighting fish (*Betta splendens* Regan, 1910) has been providing the highest income among exported ornamental fish in Thailand. During the fish life span, live diets such as rotifers, infusorians, water fleas (*Moina* sp.) and mosquito larvae are mainly used. Propagation of the live diets mostly uses the wastes from avian and porcine farms that cause the incidence of diseases and environmental impacts. These have contributed to the decrease in survival rate of juveniles and slow growth rate in maturing fish. In order to increase successive growth and survival of juvenile fish, artificial diets with improved nutrient utilization are important. Many methodologies were used to increase nutrient utilization, such as microwave cooking (Negi et al., 2001; Alajaji and El-Adawy, 2006; Khatoon and Prakash, 2006; Sadeghi and Shawrang, 2006; Hu and Wen, 2008; Ma et al., 2009), gamma irradiation (Al-Masri and Guenther, 1999; Fombang et al., 2005; El-Niely, 2007; Ebrahimi et al., 2009; Chung et al., 2010; Yoon et al., 2010), probiotics (Yanbo and Zirong, 2006; Son et al., 2009) and digestive enzymes supplementation (Mohapatra et al., 2002; Kumar et al., 2006; Lin et al., 2007). Carnivorous fishes, including Siamese fighting fish, have limited ability for carbohydrate digestion, especially at juvenile stage, due to short intestine and low activity of carbohydrate digestive enzymes. However, dietary carbohydrate appears to be necessary for improving growth and protein utilization in many fish species (Wilson, 1994). Mohapatra et al. (2002) reported a significant increase in carbohydrate utilization in *Labeo rohita* fry with increasing the level of gelatinized carbohydrate or by supplementing carbohydrases.

The objective of this study was to select a suitable modified method for improving nutrient utilization, based on equal nutritional values, in juvenile Siamese fighting fish. The main feed ingredients were selected based on *in vitro* digestibility studies of protein and carbohydrate using trypsin activity and amylase activity for standardization, respectively

(Thongprajukaew, 2011). The formulated feed was then treated by different procedures for improving nutrient utilization. Digestive enzyme specific activities and muscle compositions were used for determining physiological alterations during the experiment. Digestive enzyme indicators were measured; activity ratio of amylase to trypsin (A/T ratio) for feeding habit and metabolic flexibility of carbohydrate-protein utilizations (Hofer and Schiemer, 1981) and activity ratio of trypsin to chymotrypsin (T/C ratio) for evaluating growth efficiency (Sunde et al., 2001; Sunde et al., 2004; Rungruangsak-Torrissen, 2007). The diet with high carbohydrate content was chosen to test the hypothesis. Two week experiments were conducted at intensive rearing period after juvenile digestive tract was completely developed. These studies could provide knowledge for improving diet quality for rearing juvenile Siamese fighting fish.

2. Materials and methods

2.1. Experimental diets

2.1.1. Preliminary study

In vitro digestibility was performed for screening appropriate dose for gamma irradiation and appropriate time for microwave irradiation for modifying diets. Appropriate feedstuffs for culturing juvenile Siamese fighting fish were selected based on the *in vitro* protein and carbohydrate digestibilities, as described by Thongprajukaew (2011). The main feed mixture used for gamma or microwave irradiation contained fish meal (30%), soybean meal (20%), wheat gluten (12%), squid meal (5%) and wheat flour (20%), as shown in Table 1. For gamma irradiation, the main feed mixture was irradiated at the dose of 20, 40, 60 or 80 kGy using ^{60}Co as gamma irradiation source (Thailand Institute of Nuclear Technology, Thailand). For microwave irradiation, 100 g of the main feed mixture was placed in a plastic box (20 cm diameter \times 10 cm height), mixed with distilled water (1:4 w/v) and then cooked

at 700 W in a microwave oven (SANYO, Model EM-700T, 2450 MHz) under agitation for 4, 8, 12, 16 or 20 min. The irradiated feed mixtures were kept at 4°C until used. They were freeze-dried, using Heto FD3 (Heto-Holten, Denmark), for 2 days before *in vitro* digestibility studies of protein and carbohydrate were performed by using enzyme extracts from 20 days old juvenile Siamese fighting fish.

2.1.2. Preparation of experimental diets

The ingredients of experimental diets are shown in Table 1. The unmodified diet (control) was produced by mixing the main feed mixture with additives and vitamin-mineral premixes, and then water (30%) was added to make appropriate moisture. The glutinous mixture was passed through a hand pelletizer, then dried at 60°C for 3 h, and stored at 4°C until used. The modified diets were prepared by four different processes. 1) Gamma-irradiated diet and 2) Microwave-irradiated diet were prepared by irradiating the main feed mixture using gamma source from ^{60}Co or microwave oven, respectively, at the best dose and time obtained by highest *in vitro* digestibility values from the preliminary study in 2.1.1. The irradiated main feed mixtures were then mixed with the minor ingredients (see Table 1). 3) Probiotic-supplemented diet was freshly prepared by spraying the unmodified diet with probiotic, *Lactobacillus plantarum* KKU CRIT5 (Premier CO., LTD, Thailand) before used. The population level of *L. plantarum* in the diet was 2.7×10^8 CFU per g diet. 4) Carbohydrases-supplemented diet was prepared by spraying the unmodified diet with a mixture of carbohydrases ($100 \mu\text{l kg diet}^{-1}$), then dried at ambient temperature, and stored at 4°C until used. The mixture of the enzymes was from *Bacillus lentus* (Behn Meyer Chemical Co., Ltd., Thailand) containing the main mannan-digesting enzymes, β -mannanase, and the minor enzymes of amylase, β -glucanase, xylanase, cellulase and α -galactosidase. The

required amount of the carbohydrases was dissolved in distilled water before used. All modified diets were pelleted and kept in the same way as the control diet.

2.1.3. Biochemical composition study

The diets were dried at 105°C for 24 h before analyzing protein, lipid, fiber, and ash, as described by the AOAC (2005). The values were expressed as % on dry matter basis. Carbohydrate values or nitrogen free extract (NFE) were calculated by the difference.

2.1.4. Evaluation of gelatinization degree and water solubility

The diets were freeze-dried for two days before analysis. The dried mass of the diets was determined for degree of starch gelatinization according to Guraya and Toledo (1993). Water solubility of all nutrients was measured according to the method of Chung et al. (2010).

2.2. Fish husbandry and sample collection

Juvenile fish were obtained from a private farm in Nakhon Pathom Province, the most important area for producing exported Siamese fighting fish in Thailand. The fish were acclimatized indoors, in tanks (60 cm diameter × 30 cm height) with water temperature of $28.5 \pm 0.3^{\circ}\text{C}$, and fed with the control (unmodified) diet for 7 days before starting the experiments. The fish of 72.73 ± 2.14 mg initial weight and 19.70 ± 0.04 mm initial length were randomly distributed into 15 aquaria (18×19×34 cm), 30 fish per aquarium with a porous white cubic box (6×16×22 cm) for reducing aggressive stress between fish members. The experiment was conducted for 2 weeks with 12-h light/12-h dark and performed in triplicate with five dietary groups (one control and four modified diets) comprised of 90 fish each group. The fish were fed *ad libitum*, twice daily at 08:00 and 18:00 h. At the end of the experiment, the fish were sacrificed by chilling in ice according to “Ethical Principles and Guidelines for the Use of Animals for Scientific Purposes”, National Research Council,

Thailand. The fish were not fed on the sampling day. Body weight and total length were measured before white muscle and digestive tracts were carefully collected. The tissues were then kept at -80°C until analyses.

Weight and length of the juvenile fish were measured individually. Growth performance parameters were calculated as the following formulae.

$$\text{Condition factor (g cm}^{-3}\text{)} = 100 \times (W/L^3),$$

where W = live body weight (g) and L = total body length (cm).

Specific growth rate (SGR) was calculated according to Houde and Schekter (1981).

$$\text{SGR (\% day}^{-1}\text{)} = 100[e^g - 1]$$

where $g = (\ln W_t - \ln W_0)/(t - t_0)$, W_t = mean weight at month t , W_0 = mean weight at month t_0 .

$$\text{Net weight gain (NWG)} = \text{final body weight} - \text{initial body weight}$$

$$\text{Average daily growth (ADG, g day}^{-1}\text{)} = \text{net weight gain} / \text{rearing period}$$

$$\text{Digestosomatic index (DSI, \%)} = 100 \times [\text{gastrointestinal weight} / \text{body weight}]$$

2.3. Water quality management

The experiments were conducted at Kasetsart University in an indoor recirculating aquaculture system with a flow-rate of 280 ml min^{-1} . The recirculating aquaculture system was modified from Kovitvadhi et al. (2008). This system consisted of particulate filter cabinet ($L \times W \times H = 35 \times 22 \times 51 \text{ cm}$), macrophytes filter cabinet ($35 \times 85 \times 51 \text{ cm}$), biological filter cabinet ($35 \times 72 \times 51 \text{ cm}$), water resting cabinet ($35 \times 35 \times 51 \text{ cm}$), and five culture units ($34 \times 19 \times 26 \text{ cm}$). All cabinets had the water level of 45 cm while the culture units had the water level of 18 cm. The water parameters were analyzed twice weekly. Water temperature, pH, conductivity and dissolved oxygen were analyzed using water analyzer (Multi probe system, 556 MPS, YSI Incorporated, USA). Other parameters including total alkalinity (phenolphthalein methyl orange indicator), free carbon dioxide (titration), total hardness (EDTA titration), total ammonia nitrogen (phenate method), nitrite (colorimetry), nitrate

(cadmium reduction) and orthophosphate (ascorbic acid method) were analyzed according to the method of APHA, AWWA, WPCF (1998).

The water quality during the experimental period had the temperature of 28.49 ± 0.28 °C, pH 7.52 ± 0.05 , dissolved oxygen 3.95 ± 0.06 mg L⁻¹, conductivity 0.40 ± 0.01 mS cm⁻¹, total alkalinity 94.74 ± 1.13 ppm CaCO₃, total hardness 114.75 ± 0.81 ppm CaCO₃, free carbon dioxide 1.38 ± 0.05 ppm, nitrate 0.045 ± 0.003 ppm, nitrite 0.0033 ± 0.0001 ppm, total ammonia nitrogen 0.027 ± 0.004 ppm, and phosphorous 0.028 ± 0.002 ppm.

2.4. Digestive enzyme studies

2.4.1. Enzyme extraction

The enzyme extractions were performed according to Rungruangsak-Torrissen (2007). Digestive tracts of juvenile fish were extracted in 50 mM Tris-HCl buffer pH 8 containing 200 mM NaCl (1:3 w/v) using micro-homogenizer (THP-220, OMNI International, USA). The homogenate was centrifuged at $10,000 \times g$ for 20 min at 4°C. The supernatant was then collected and kept at -80°C in small portions for later determinations. Protein concentration in the crude enzyme extract was determined according to Lowry et al. (1951).

2.4.2. Digestive enzyme assays

The optimal conditions (pH and temperature) chosen for studying the main digestive enzymes in Siamese fighting fish were according to Thongprajukaew et al. (2010a, 2010b).

Amylase activity (at pH 8 and 50°C) was determined based on Areekijseree et al. (2004) modified from Bernfeld (1951) using starch solution as substrate. The enzyme digestion reaction was modified to 15 min. Amylase specific activity was expressed as μmol maltose produced h⁻¹ mg protein⁻¹.

Total protease activity (at pH 8 and 50°C) was assayed using azocasein as substrate based on Areekijserree et al. (2004) modified from Garcia-Carreno (1992). The specific activity of total protease was expressed as mU mg protein⁻¹. One unit (U) of total protease activity was defined as the amount of enzyme giving an increase of 1.0 absorbance unit at 440 nm at the specified reaction condition.

Amidase activities of trypsin (at pH 8 and 50°C) and chymotrypsin (at pH 8 and 50°C) were assayed by initial reactions based on Rungruangsak-Torrissen (2007) using BAPNA (benzoyl-*L*-arginine-*p*-nitroanilide) and SAPNA (*N*-succinyl-ala-ala-pro-phe-*p*-nitroanilide) as specific substrates, respectively. The specific activities of trypsin and chymotrypsin were expressed as $\mu\text{mol } p\text{-nitroaniline produced h}^{-1} \text{ mg protein}^{-1}$.

Esterase activity of lipase (at pH 8 and 40°C) was analyzed based on Winkler and Stuckmann (1979) using *p*-nitrophenyl palmitate as substrate. The specific activity of lipase was expressed as $\mu\text{mol } p\text{-nitrophenol produced h}^{-1} \text{ mg protein}^{-1}$.

2.5. *In vitro* digestibility studies

Crude enzyme extracts were dialyzed overnight against 50 mM Tris-HCl buffer pH 8.2 before used for determining *in vitro* digestibility. Freeze-dried diets were used as substrate. Protein and carbohydrate digestibilities of the experimental diets using fish crude enzyme extracts were determined using the method modified from Rungruangsak-Torrissen et al. (2002) and Areekijserree et al. (2006). The reaction mixture containing 5 mg dried feed, 10 ml 50 mM phosphate buffer pH 8.2, 50 μl 0.5 % chloramphenical, and 125 μl dialyzed crude enzyme extract, was incubated at 25°C for 24 h.

Protein digestibility was determined by measuring the increase in liberated reactive amino groups of cleaved peptides. The reaction mixture, containing 200 μl digested solution, 2 ml 50 mM phosphate buffer pH 8.2, and 1 ml 0.1% trinitrobenzene sulphonic acid (TNBS),

was heated in the dark at 60 °C for 1 h, and stopped by adding 1 ml 1 M HCl before measuring absorbance at 420 nm and comparison with *DL*-alanine standard curve.

Carbohydrate digestibility was determined by measuring the increase in reducing sugar. The reaction mixture containing 1 ml digested solution and 500 µl DNS, was heated in boiling water for 5 min and cooled to room temperature before measuring absorbance at 540 nm and comparison with maltose standard curve.

The blanks (without dialyzed crude enzyme extracts) were used to deduct liberated amino acids and reducing sugars. For comparison, the calculated values were standardized by trypsin activity for protein digestibility and by amylase activity for carbohydrate digestibility (Thongprajukaew, 2011). The *in vitro* digestibility of protein was expressed as µmol *DL*-alanine equivalent g dried feed⁻¹ trypsin activity⁻¹. The *in vitro* digestibility of carbohydrate was expressed as µmol maltose g dried feed⁻¹ amylase activity⁻¹.

2.6. White muscle and body compositions

Scale and skin of the fish were carefully removed, and the epaxial white muscle was dissected. RNA and protein concentrations in the muscle and body were determined as described in Rungruangsak-Torrissen (2007) modified from Sunde et al. (2001). The extinction coefficient for RNA is $E_{260} = 40 \text{ } \mu\text{g RNA ml}^{-1}$, and for protein is $E_{280} = 2.1 \text{ mg protein ml}^{-1}$. Lipids were extracted using ethyl acetate as described by Supannapong et al. (2008) and Rungruangsak-Torrissen et al. (2009). All values were expressed on wet weight basis.

2.7. Statistical analysis

Data were expressed as mean \pm standard error of mean in triplicate observations. One-Way Analysis of Variance was used for evaluating growth performance parameters, digestive

enzyme specific activities, muscle compositions and body compositions. Significant differences between means were ranked using Duncan's multiple range test (DMRT) at 95% significance level. Pearson correlation coefficients (r) between the parameters were calculated.

3. Results

3.1. Preliminary study for screening irradiation conditions

In vitro digestibilities of protein and carbohydrate in the main feed mixtures treated with different irradiation procedures are shown in Fig.1. Protein digestibility values between treated and untreated feed mixtures were not different ($P > 0.05$). The value was relatively highest in microwave irradiation for 8 min cooking time. Carbohydrate digestibility values, on the other hand, showed some differences between the modified procedures ($P < 0.05$), with highest value also in 8 min microwave cooking time. Among gamma irradiation doses, carbohydrate digestibility value was relatively highest at 20 kGy. Therefore, the 8 min microwave cooking and 20 kGy gamma irradiation were chosen as the appropriate doses for modifying diets by the two irradiation techniques.

3.2. Biochemical compositions and some physical properties of experimental diets

No differences were observed in proximate compositions and gross energy among the experimental diets (Table 2). All modified diets showed relatively higher values of *in vitro* digestibilities and physical properties than the control (Table 2). Gamma irradiation increased protein digestibility significantly compared to the control ($P < 0.05$), but the increase was not different from the other techniques ($P > 0.05$). Microwave irradiation, on the other hand, increased carbohydrate digestibility (albeit insignificant), with significantly increased degree

of starch gelatinization and water solubility ($P < 0.05$), compared to the control and the other techniques.

3.3. Survival rate and growth performance of juveniles

No significant differences ($P > 0.05$) were found in survival rate of the juveniles from all dietary treatments (Fig. 2 and Table 3). Nevertheless, the levels of fish survival were relatively higher in fish fed the experimental diets (pooled data) when compared with the control ($P > 0.05$). Survival rate of juvenile Siamese fighting fish in this experiment was ranged from 72–79%.

The results of growth performance are shown in Table 3. No significant differences were observed in total fish length between the dietary groups ($P > 0.05$). On the other hand, body weight and its related parameters were different between the dietary groups ($P < 0.05$). At the end of the experiment, fish fed on microwave cooking diet showed highest values in body weight, condition factor, specific growth rate (SGR), net weight gain (NWG), average daily gain (ADG), and gastrointestinal weight. Among the dietary groups, the levels of these parameters were observed as microwave-irradiated dietary group > carbohydrases-supplemented dietary group > gamma-irradiated dietary group > control group > probiotic-supplemented dietary group. The probiotic-supplemented dietary group also showed significantly lowest values of the digestosomatic index (DSI) ($P < 0.05$).

There was a relationship between body weight and gastrointestinal weight ($r = 0.824$, $P < 0.0001$), regardless of dietary groups (Fig. 3).

3.4. Digestive enzyme specific activities

The results of digestive enzyme specific activities are illustrated in Table 4. Amylase and total protease specific activities were different between fish groups; showing highest

levels in microwave-irradiated dietary group and lowest levels in carbohydrases-supplemented dietary group. Gamma irradiation significantly decreased total protease specific activity ($P < 0.05$), and carbohydrases supplementation significantly decreased specific activities of both amylase and total protease ($P < 0.05$), compared to the control. No differences were observed in the levels of trypsin, chymotrypsin, activity ratio of trypsin to chymotrypsin (T/C ratio), and lipase. However, the highest levels of these enzyme parameters were also observed in microwave-irradiated dietary group. These resulted in highest T/C ratio (albeit insignificant) and activity ratio of amylase to trypsin (A/T ratio, $P < 0.05$) in fish fed microwave-irradiated diet. No differences were observed in A/T ratios among the other fish groups ($P > 0.05$).

The relationships between digestive enzymes specific activities and growth, regardless of dietary groups, are shown in Table 5. Amylase specific activity of juveniles showed positive relationship with specific activities of total protease and trypsin, as well as T/C ratio. Total protease specific activity correlated with trypsin specific activity, and as usual, specific activities of trypsin and chymotrypsin are correlated. The A/T ratio correlated with amylase and total protease specific activities, as well as fish SGR. The T/C ratio correlated with trypsin specific activity, but not with chymotrypsin specific activity.

3.5. Muscle and body compositions

Muscle and body compositions of juveniles were mainly similar among fish groups, except for the levels of protein (Table 6). The levels of protein concentrations in either muscle or body were highest in the control group and lowest in the probiotic-supplemented dietary group. Muscle protein concentrations were significantly lower in carbohydrases- and probiotic-supplemented dietary groups, compared to the control ($P < 0.05$). However, their

body protein concentrations were significantly lower than the control and microwave-irradiated dietary group ($P < 0.05$).

There were no correlations between muscle or body compositions and other parameters (growth and digestive enzymes) during 2 weeks experiment.

4. Discussion

4.1. Digestibility of irradiated-main feed mixture and irradiated-diet

Differences in gamma irradiation doses and microwave cooking times did not seem to affect protein digestibility of the main feed mixture in the preliminary study, but could possibly influence carbohydrate digestibility (Fig. 1). Effects of gamma irradiation on protein digestibility have been shown to be depended on irradiation dose and type of material (Fombang et al., 2005). High correlation coefficient between radiation doses (5, 7.5 and 10 kGy) and *in vitro* digestibility of proteins from peas, cowpeas, lentil, kidney bean and chickpea has been reported (El-Niely, 2007). Higher doses of gamma irradiation tended to decrease carbohydrate digestibility, but the observations were not statistically different ($P > 0.05$, Fig. 1). This might be due to total carbohydrate digestibility was presented, as significant changes in carbohydrate digestibility have been shown in terms of starch digestible rate (Yoon et al., 2010; Chung et al., 2010). Moreover, raw materials from most studies were starch sources while the main feed mixture in this study were comprised of both protein and carbohydrate from plants and animals. For microwave cooking, its use for improving protein and carbohydrate digestibilities has been reported in various legume seeds, such as moth bean (Negi et al., 2001), green gram, Bengal gram and horse gram (Khatoon and Prakash, 2006). Intensity and irradiation time of microwave process and material concentration were main factors governing enzymatic hydrolysis in rice straw (Ma et al., 2009). This is in agreement with the observation at a proper cooking time (Fig. 1).

Microwave irradiation affected protein degradation (Sadeghi and Shawrang, 2006), which was also observed in our study using electrophoresis (SDS-PAGE) technique (results not shown).

Only gamma irradiation gave a significant increase in protein digestibility compared to the control (Table 2). The increase might be influenced by the breaking of disulphide bonds in protein molecules, as the digestibility levels were related positively with free sulphydryl group levels and negatively with disulphide bond levels (Rungruangsak-Torrissen et al., 2002). Higher quality feeds had higher levels of free sulphydryl group affected by different processing conditions (Sunde et al., 2004). Increased protein digestion in seeds by gamma irradiation was occurred by protein subunit degradation (Ebrahimi et al., 2009) and change in cell wall constituents of some agricultural by-products (Al-Masri and Guenther, 1999), allowing better contact to proteolytic enzymes. The use of different techniques also increased carbohydrate digestibility (albeit insignificant), with the highest digestibility value by microwave irradiation (Table 2). Microwave processing has been reported to improve carbohydrate digestibility of moth bean (Negi et al., 2001) and chickpea (Alajaji et al., 2006). Fish fed on gelatinized corn based diet with different levels of α -amylase supplementation showed significant increases in dry matter digestibility (Kumar et al., 2006). Increased carbohydrate digestibility was associated with increased physicochemical properties of the diets, as also observed in microwave-irradiated diet showing higher starch gelatinization and water solubility than the other diets, without affecting proximate compositions and gross energy (Table 2).

4.2. Survival rate and growth performance of juveniles

Survivals of juvenile Siamese fighting fish at the end of experiment were similar among the dietary groups (Table 3). Generally, microwave irradiation is better than the other

techniques studied, as it improved physicochemical properties of the diets and growth of the juveniles (Tables 2 and 3). The diet may be more palatable than the other diets, as the fish were observed to take shorter time to ingest the microwave-irradiated diet. Carbohydrases supplementation did not improve fish growth performance (Table 3; Rungruangsak-Torrissen et al., 2010), similar to the observations of supplementations with hemicellulose digesting enzymes and α -galactosidase in lupin-based diets (Lin et al., 2007). Uses of probiotics for enhancing successive growth have been reported by Yanbo and Zirong (2006) and Son et al. (2009), however, it did not improve growth performance in our experiment (Table 3). Gastrointestinal weight and digestosomatic index (DSI) of juvenile fish fed on the modified diets were mainly similar to those of the control fish, except for supplementation with probiotic *L. plantarum* resulted in decreased DSI (Table 3). A decrease in the index was also reported in juvenile rohu (*Labeo rohita*) feeding on gelatinized corn based diet with or without α -amylase supplementation (Kumar et al., 2006).

4.3. Responses of digestive enzymes

The increased gelatinization and water solubility of the microwave-irradiated diet could have affected on the higher digestive enzyme specific activities, which contributed to the higher growth performance in this fish group (Tables 2–4). Increased gelatinization level of dietary carbohydrate resulting in increased amylase activity was also observed in rahu, *Labeo rohita* (Mohapatra et al., 2002). Up-regulations of enzymes involved in carbohydrate and protein digestions in microwave-irradiated dietary group could be due to greater utilization of both nutrients for higher energy requirement and growth performance, without changing muscle and body compositions (Table 6). However, down-regulated of these enzymes in carbohydrases-supplemented dietary group caused significantly lower protein levels in both body and muscle (Tables 4 and 6), because of lower in both energy and protein

utilizations than the control. Changes in dietary carbohydrate and protein structures (through microwave or gamma irradiations) and supplementations of exogenous enzymes (from *L. plantarum* or *B. lentus*), could influence endogenous enzymes productions for utilizing energy and nutrients for fish growth performance. These are supported by the relationships observed between the levels of amylase, total protease, trypsin, T/C ratio and A/T ratio (Table 5). These parameters were higher in higher growth group (microwave-irradiated dietary group), although some parameters may not show statistically different (Tables 3 and 4). The specific activities of the alkaline proteases trypsin and chymotrypsin, including T/C ratio, are important for understanding growth performance quality and feed utilization efficiency in different fish species (Sunde et al., 2001, 2004; Rungruangsak-Torrissen, 2007; Rungruangsak-Torrissen and Fosseidengen, 2007; Rungruangsak-Torrissen et al., 2009, 2010) including Siamese fighting fish, however, the A/T ratio was not found to associate with fish growth (Thongprajukaew, 2011). The significantly higher growth performance and A/T ratio, but not T/C ratio, in microwave-irradiated dietary group may indicate higher energy requirement for protein utilization and growth in this group than the other groups. Lipid utilization of the fish did not seem to be affected by the modified diets, as the specific activity of lipase and lipid deposition in body and muscle were similar to the control (Tables 4 and 6).

4.4. Muscle and carcass of juveniles

Similarity of muscle and carcass compositions in microwave- and gamma- irradiated dietary groups, compared to control group, indicated better growth of juvenile Siamese fighting fish feeding on these irradiated diets. The uses of diets with probiotic- and carbohydrases- supplementations, on the other hand, reduced protein depositions in body and muscle of the fish, compared to the control. The modified diets did not affect capacities for protein synthesis (RNA concentration) and turnover (RNA/Protein ratio), and protein growth

(Protein/Lipid ratio) in Siamese fighting fish. The experimental period of 2 weeks might be too short to observe the differences. However, 10 weeks feeding on pre-gelatinized starch had no effect on whole body and muscle compositions in juvenile European sea bass, *Dicentrarchus labrax* (Peres and Oliva-Teles, 2002) and juvenile yellowfin seabream (Wu et al., 2007). Increased *in vitro* protein digestibility of diet with fish meal as the whole protein source could increase feed efficiency and muscle protein synthesis capacity in fish (Rungruangsak-Torrissen et al., 2002; Sunde et al., 2004). On the other hand, increased *in vitro* protein digestibility of diets with high plant protein did not improve fish growth performance quality as well as diets with high animal protein did (Rungruangsak-Torrissen et al., 2010). The lower protein depositions in probiotic- and carbohydrases- supplemented dietary groups, and the lack of increased protein depositions in fish fed gamma-irradiated diet with increased *in vitro* protein digestibility value, compared to control, might be due to a high level of dietary plant proteins that could cause imbalance of amino acids in these diets.

5. Conclusion

Modifications of diets by irradiations (with 20 kGy gamma ray from ^{60}Co or 8 min microwave cooking at 700 W) or supplementations (with probiotic *L. plantarum* 2.7×10^8 CFU per g diet or enzymes mixture of commercial carbohydrases 100 μl per kg diet) could affect qualities of dietary nutrients. Gamma-irradiated diet showed higher *in vitro* protein digestibility than control diet, and it decreased total protease specific activity without changing fish growth performance quality. Microwave-irradiated diet, with insignificant improved *in vitro* digestibilities of protein and carbohydrate, showed significant improved starch gelatinization and water solubility that could improve fish growth performance through increased amylase specific activity and A/T ratio without changing muscle and body compositions of the fish. Probiotic-supplemented diet, with similar dietary qualities that

resulted in similar levels of digestive enzymes and fish growth as the control diet, decreased protein depositions in body and muscle of the fish. Carbohydrases-supplemented diet, with similar dietary qualities that resulted in similar fish growth as the control diet, decreased the levels of amylase and total protease including protein depositions in body and muscle of the fish. The modified diets did not show any effect on lipid utilization of the fish. There were interactions between carbohydrate and protein utilizations, as shown by the relationships among the levels of amylase, total protease, trypsin, T/C ratio and A/T ratio. All fish groups showed similar capacity for protein synthesis and turnover including protein growth. Changes in digestive enzymes specific activities and protein depositions in some dietary groups should probably be due to consumption rate as the fish were fed *ad libitum*. Enhanced growth performance in microwave-irradiated dietary group may not only be due to improved nutrient quality in the diet but also improved fish consumption rate.

Acknowledgements

We would like to thank Prof. Dr. Busaba Yongsmith, Department of Microbiology, Kasetsart University, for kindly preparing red monascol rice; and Premier CO., LTD., Thailand, for kindly providing the probiotic. This work was financially supported by the Office of the Higher Education Commission (OHEC), Thailand, for the research funding under the program Strategic Scholarships for Frontier Research Network for the Joint Ph.D. Program Thai Doctoral Degree; and in part by Thesis and Dissertation Research Fund, the Graduate School, Kasetsart University, Thailand.

References

- 467 Al-Masri, M. R., Guenther, K. D., 1999. Changes in digestibility and cell-wall constituents of
468 some agricultural by-products due to gamma irradiation and urea treatments. *Radiat.*
469 *Phys. Chem.* 55, 323–329.
- 470 Alajaji, S. A., El-Adawy, T. A., 2006. Nutritional composition of chickpea (*Cicer arietinum*
471 L.) as affected by microwave cooking and other traditional cooking methods. *J. Food*
472 *Comp. Anal.* 19, 806–812.
- 473 AOAC, 2005. Official Methods of Analysis of AOAC International, 18th ed. Association of
474 Official Analytical Chemists, Maryland, USA.
- 475 APHA, AWWA, WPCF, 1998. Standard methods for the examination of water
476 and wastewater, American Public Health Association, American Water Works
477 Association, 20th ed. Water Pollution Control Federation, Washington, DC.
- 478 Areekijseeree, M., Engkagul, A., Kovitvadhi, U., Thongpan, A., Mingmuang, M., Pakkong, P.,
479 Rungruangsak-Torrissen, K., 2004. Temperature and pH characteristics of amylase and
480 proteinase of adult freshwater pearl mussel, *Hyriopsis (Hyriopsis) bialatus* Simpson
481 1900. *Aquaculture* 234, 575–587.
- 482 Areekijseeree, M., Engkagul, A., Kovitvadhi, S., Kovitvadhi, U., Thongpan, A.,
483 Rungruangsak-Torrissen, K., 2006. Development of digestive enzymes and *in vitro*
484 digestibility of different species of phytoplankton for culture of early juveniles of the
485 freshwater pearl mussel, *Hyriopsis (Hyriopsis) bialatus* Simpson, 1900. *Invert. Reprod.*
486 *Dev.* 49, 255–262.
- 487 Bernfeld, P., 1951. Enzymes of starch degradation and synthesis. *Adv. Enzymol.* 12, 379.
- 488 Chung, H-J., Lee, S-Y., Kim, J-H., Lee, J-W., Byun, M-W., Lim, S-T., 2010. Pasting
489 characteristics and *in vitro* digestibility of γ -irradiated RS₄ waxy maize starches. *J.*
490 *Cereal Sci.* 52, 53–58.

- 491 Ebrahimi, S. R., Nikkhah, A., Sadeghi, A. A., Raisali, G., 2009. Chemical composition,
492 secondary compounds, ruminal degradation and *in vitro* crude protein digestibility of
493 gamma irradiated canola seed. Anim. Feed Sci. Technol. 151, 184–193.
- 494 El-Niely, H. F. G., 2007. Effect of radiation processing on antinutrients, *in vitro* protein
495 digestibility and protein efficiency ratio bioassay of legume seeds. Radiat. Phys. Chem.
496 76, 1050–1057.
- 497 Fombang, E. N., Taylor, J. R. N., Mbofung, C. M. F., Minnaar, A., 2005. Use of γ -irradiation
498 to alleviate the poor protein digestibility of sorghum porridge. Food Chem. 91, 695–703.
- 499 Garcia-Carreno, F.L., 1992. The digestive proteases of langostilla (*Pleuroncodes palanipes*,
500 Decapoda): their partial characterization and the effect of food on their composition.
501 Comp. Biochem. Physiol. 103B, 575–578.
- 502 Guraya, H. S., Toledo, R. T., 1993. Determining gelatinized starch in a dry starchy product. J.
503 Food Sci. 58, 888.
- 504 Hofer, R., Schiemer, F., 1981. Proteolytic activity in the digestive tract of several species of
505 fish with different feeding habits. Oecologia 48, 342–345.
- 506 Houde, E.D., Schekter, R.C., 1981. Growth rate, rations and cohort consumption of marine
507 fish larvae in relation to prey concentration. Rapp. P-V Reun. Cons. Int. Explor. Mer
508 178, 441–453.
- 509 Hu, Z., Wen, Z., 2008. Enhancing enzymatic digestibility of switchgrass by microwave-
510 assisted alkali pretreatment. Biochem. Eng. J. 38, 369–378.
- 511 Khatoon, N., Prakash, J., 2006. Nutrient retention in microwave cooked germinated legumes.
512 Food Chem. 97, 115–121.
- 513 Kovitvadhi S., Kovitvadhi U., Sawangwong P., Machado J., 2008. A laboratory-scale
514 recirculating aquaculture system for juveniles of freshwater pearl mussel *Hyriopsis*
515 (*Limnoscapha*) *myersiana* (Lea, 1856). Aquaculture 49, 255-262.

- 516 Kumar, S., Sahu, N. P., Pal, A. K., Choudhury, D., Mukherjee, S. C., 2006. Studies on
517 digestibility and digestive enzyme activities in *Labeo rohita* (Hamilton) juveniles:
518 effect of microbial α -amylase supplementation in non-gelatinized or gelatinized corn-
519 based diet at two protein levels. Fish Physiol. Biochem. 32, 209–220.
- 520 Lin, S., Mai, K., Tan, B., 2007. Effects of exogenous enzyme supplementation in diets on
521 growth and feed utilization in tilapia, *Oreochromis niloticus* \times *O. aureus*. Aquacult. Res.
522 38, 1645–1653.
- 523 Lowry, O. H., Rosenbrough, N. J., Farr, A. L., Randall, R. J., 1951. Protein measurement
524 with Folin phenol reagent. J. Biol. Chem. 193, 265–275.
- 525 Ma, H., Liu, W-W., Chen, X., Wu, Y-J., Yu, Z-L., 2009. Enhanced enzymatic
526 saccharification of rice straw by microwave pretreatment. Biores. Technol. 100, 1279–
527 1284.
- 528 Mohapatra, M., Sahu, N. P., Chaudhari, A., 2002. Utilization of gelatinized carbohydrate in
529 diets in *Labeo rohita* fry. Aquacult. Nutr. 8, 1–8.
- 530 Negi, A., Boora, P., Khetarpaul, N., 2001. Effect of microwave cooking on the starch and
531 protein digestibility of some newly released moth bean (*Phaseolus aconitifolius* Jacq.)
532 cultivars. J. Food Comp. Anal. 14, 541–546.
- 533 Peres, H., Oliva-Teles, A., 2002. Utilization of raw and gelatinized starch by European sea
534 bass (*Dicentrarchus labrax*) juveniles. Aquaculture 205, 287– 299.
- 535 Rungruangsak-Torrissen, K., 2007. Digestive efficiency, growth and qualities of muscle and
536 oocyte in Atlantic salmon (*Salmo salar* L.) fed on diets with krill meal as an alternative
537 protein source. J. Food Biochem. 31, 509–540.
- 538 Rungruangsak-Torrissen, K., Rustad, A., Sunde, J., Eiane, S. A., Jensen, H. B., Opstvedt, J.,
539 Nygard, E., Samuelsen, T. A., Mundheim, H., Luzzana, U., Venturini, G., 2002. *In vitro*

- 540 digestibility based on fish crude enzyme extract for prediction of feed quality in growth
541 trials. J. Sci. Food Agric. 82, 644–654.
- 542 Rungruangsak-Torrissen, K., Fosseidengen, J. E., 2007. Effect of artificial feeding on
543 digestive efficiency, growth and qualities of muscle and oocyte of maturing Atlantic
544 mackerel (*Scomber scombrus* L.). J. Food Biochem. 31, 726–747.
- 545 Rungruangsak-Torrissen, K., Kovitvadhi, U., Siruntawinetti, J., Trenet, E., Engkagul, A.,
546 Aidnoie, Y., Choowongkamon, K., Unajak, S., Meeswad, P., Sunthornchot, J.,
547 Kovitvadhi, S., Preprame, P., 2010. Development of suitable feed for commercial
548 production of Nile tilapia, *Oreochromis niloticus*. Kasetsart University Technical
549 Report 2010, 184 pp. (in Thai). Biochemical Research Unit for Feed Utilization
550 Assessment, Kasetsart University, Bangkok, Thailand.
- 551 Rungruangsak-Torrissen, K., Stien, L.H., Daae, B.S., Vågseth, T., Thorsheim, G.B., Tobin,
552 D., Ritola, O., 2009. Different dietary levels of protein to lipid ratio affected digestive
553 efficiency, skeletal growth, and muscle protein in rainbow trout families. Scholarly
554 Research Exchange, vol. 2009, Article ID 709529, doi: 10.3814/2009/709529.
- 555 Sadeghi, A. A., Shawrang, P., 2006. Effects of microwave irradiation on ruminal
556 degradability and *in vitro* digestibility of canola meal. Anim. Feed Sci. Technol. 127,
557 45–54.
- 558 Son, V. M., Chang, C-C., Wu, M-C., Guu, Y-K., Chiu, C-H., Cheng, W., 2009. Dietary
559 administration of the probiotic, *Lactobacillus plantarum*, enhanced the growth, innate
560 immune responses, and disease resistance of the grouper *Epinephelus coioides*. Fish
561 Shell. Immunol. 26, 691–698.
- 562 Sunde, J., Eiane, S. A., Rustad, A., Jensen, H. B., Opstvedt, J., Nygard, E., Venturini, G.,
563 Rungruangsak-Torrissen, K., 2004. Effect of fish feed processing conditions on

- 564 digestive protease activities, free amino acid pools, feed conversion efficiency and
 565 growth in Atlantic salmon (*Salmo salar* L.). *Aquacult. Nutr.* 10, 261–277.
- 566 Sunde, J., Taranger, G.L., Rungruangsak-Torrissen, K., 2001. Digestive protease activities
 567 and free amino acids in white muscle as indicators for feed conversion efficiency and
 568 growth rate in Atlantic salmon (*Salmo salar* L.). *Fish Physiol. Biochem.* 25, 335–345.
- 569 Supannapong, P., Pimsalee, T., A-komol, T., Engkagul, A., Kovitvadhi, U., Kovitvadhi, S.,
 570 Rungruangsak-Torrissen, K., 2008. Digestive enzymes and *in vitro* digestibility of
 571 different species of phytoplankton for culture of the freshwater pearl mussel, *Hyriopsis*
 572 (*Hyriopsis*) *bialatus*. *Aquacult. Int.* 16, 437–453.
- 573 Thongprajukaew, K., 2011. Feed development using digestive enzyme technology for
 574 successive growth in Siamese fighting fish (*Betta splendens* Regan, 1910). Ph.D. thesis,
 575 139 pp. Kasetsart University, Inter Departmental Multidisciplinary Graduate Program in
 576 Bioscience, Bangkok, Thailand.
- 577 Thongprajukaew, K., Kovitvadhi, U., Engkagul, A., Rungruangsak-Torrissen, K. 2010a.
 578 Characterization and expression levels of protease enzymes at different developmental
 579 stages of Siamese fighting fish (*Betta splendens* Regan, 1910). *Kasetsart J. (Nat. Sci.)*
 580 44 (3), 411–423.
- 581 Thongprajukaew, K., Kovitvadhi, U., Engkagul, A., Rungruangsak-Torrissen, K., 2010b.
 582 Temperature and pH characteristics of amylase and lipase at different developmental
 583 stages of Siamese fighting fish (*Betta splendens* Regan, 1910). *Kasetsart J. (Nat. Sci.)*
 584 44 (2), 210–219.
- 585 Wilson, R. P., 1994. Utilization of dietary carbohydrate by fish. *Aquaculture* 124, 67–80.
- 586 Winkler, U. K., Stuckmann, M., 1979. Glycogen, hyaluronate and some other
 587 polysaccharides greatly enhance the formation of exolipase by *Serratia marcescens*. *J.*
 588 *Bacteriol.* 138, 663–670.

- 589 Wu, X-Y., Liu, Y-J., Tian, L-X., Mai, K-S., Gou, R., Jin, S-J., 2007. Effect of different
590 dietary raw to pre-gelatinized starch ratios on growth performance, feed utilization and
591 body composition of juvenile yellowfin seabream (*Sparus latus*). Aquacult. Int. 15,
592 467–477.
- 593 Yanbo, W., Zirong, X., 2006. Effect of probiotics for common carp (*Cyprinus carpio*) based
594 on growth performance and digestive enzyme activities. Anim. Feed Sci. Technol. 127,
595 283–292.
- 596 Yoon, H-S., Yoo, J-Y., Kim, J-H., Lee, J-W., Byun, M-W., Baik, B-K., Lim, S-T., 2010. *In*
597 *vitro* digestibility of gamma-irradiated corn starches. Carbohydr. Pol. 81, 961–963.

Figure captions

Fig. 1 *In vitro* digestibilities of protein ($\mu\text{mol } DL\text{-alanine equivalent g dried feed}^{-1}$ trypsin activity $^{-1}$) and carbohydrate ($\mu\text{mol maltose g dried feed}^{-1}$ amylase activity $^{-1}$) of unmodified, gamma-irradiated (20–80 kGy) and microwave-irradiated (700 W for 4–20 min) main feed mixtures, using dialyzed crude enzyme extracts from 20-day-old Siamese fighting fish. Data was calculated from triplicate observations. The values with different superscripts are significantly different ($P < 0.05$).

Fig. 2 Change in survival rate (%) of juvenile Siamese fighting fish fed the different diets. Data was expressed as the mean of triplicate aquaria. There was no significant difference ($P > 0.05$) among the dietary groups at the end of rearing period.

Fig. 3 General relationship between body weight and gastrointestinal weight of juvenile Siamese fighting fish in all dietary groups.


Fig. 1


Fig. 2


Fig. 3

Table 1

Ingredients of the experimental diets for rearing juvenile Siamese fighting fish.

Ingredients	Inclusion (%)
Fish meal*	30
Soybean meal*	20
Wheat gluten*	12
Squid meal	5
Wheat flour*	20
Lecithin	2
Fish oil	1
Soybean oil	2.6
Mineral mixture**	0.05
Vitamin mixture***	0.25
Vitamin C	0.1
Fermented red rice	2
Cellulose	5

* Main ingredients selected from *in vitro* screening of suitable feedstuffs using the enzyme extracts from juvenile Siamese fighting fish, as described by Thongprajukaew (2011).

** Mineral mixtures, 1 kg of feed contained 30 mg iron, 20 mg zinc, 25 mg manganese, 5 mg copper, 5 mg iodine and 0.2 mg selenium.

*** Vitamin mixtures, 1 kg of feed contained 4,000 IU vitamin A, 2,000 IU vitamin D₃, 50 mg vitamin E, 10 mg vitamin K, 20 mg thiamine, 20 mg riboflavin, 20 mg pyridoxine, 200 mg calcium panthothenate, 150 mg niacin, 2 mg biotin, 5 mg folic acid, 0.2 mg vitamin B₁₂, 400 mg inositol and 200 mg ethoxyquin.

Table 2

Biochemical compositions (on dry matter basis), *in vitro* digestibilities of protein ($\mu\text{mol } DL\text{-alanine equivalent g dried feed}^{-1} \text{ trypsin activity}^{-1}$) and carbohydrate ($\mu\text{mol maltose g dried feed}^{-1} \text{ amylase activity}^{-1}$), starch gelatinization (%), and water solubility (%) of the experimental diets used for rearing juvenile Siamese fighting fish. Data were obtained from triplicate observations.

Dietary parameters	Experimental diets				
	Unmodified	Gamma irradiation	Microwave irradiation	Probiotic	Carbohydrases
Crude protein (%)	41.9	41.8	42.3	42.4	42.6
Crude lipid (%)	7.0	7.2	7.0	7.0	6.8
Nitrogen free extract (%)	32.9	32.1	32.0	31.6	31.1
Crude fiber (%)	4.1	4.5	4.0	4.5	4.9
Ash (%)	14.1	14.4	14.7	14.5	14.6
Gross energy (kJ g^{-1})	19.5	19.4	19.3	19.3	19.2
<i>In vitro</i> protein digestibility	200.23 ± 9.80^b	234.78 ± 8.88^a	221.24 ± 10.77^{ab}	216.26 ± 8.07^{ab}	222.61 ± 8.93^{ab}
<i>In vitro</i> carbohydrate digestibility	1.80 ± 0.01	1.82 ± 0.17	2.48 ± 0.20	2.13 ± 0.47	1.92 ± 0.11
Starch gelatinization	63.69 ± 0.92^b	66.34 ± 0.35^b	76.23 ± 0.13^a	64.01 ± 0.83^b	63.89 ± 0.41^b
Water solubility	19.32 ± 0.26^b	22.23 ± 0.25^b	30.76 ± 0.63^a	20.02 ± 1.02^b	19.97 ± 0.71^b

The values in the same row with different superscripts are significantly different ($P < 0.05$).

Table 3

Effects of the experimental diets on survival rate and growth performance of juvenile Siamese fighting fish at the end of experiment. Data were obtained from triplicate observations.

Parameters	Unmodified	Gamma irradiation	Microwave irradiation	Probiotic	Carbohydases
Survival rate (%)	72.22 ± 2.22	78.89 ± 1.11	77.78 ± 2.94	75.56 ± 2.94	76.67 ± 6.67
Total length (mm)	22.08 ± 0.14	22.33 ± 0.46	22.88 ± 0.40	21.81 ± 0.23	22.40 ± 0.34
Body weight (mg)	111.72 ± 6.88 ^b	118.13 ± 6.61 ^{ab}	134.75 ± 1.03 ^a	108.00 ± 5.15 ^b	123.28 ± 8.61 ^{ab}
Condition factor (g cm ⁻³)	1.03 ± 0.03 ^b	1.04 ± 0.02 ^b	1.15 ± 0.03 ^a	1.01 ± 0.01 ^b	1.06 ± 0.05 ^{ab}
Specific growth rate (% day ⁻¹)	3.09 ± 0.45 ^b	3.50 ± 0.41 ^{ab}	4.50 ± 0.06 ^a	2.85 ± 0.36 ^b	3.81 ± 0.51 ^{ab}
Net weight gain (mg)	38.98 ± 6.88 ^b	45.40 ± 6.61 ^{ab}	62.02 ± 1.03 ^a	35.27 ± 5.15 ^b	50.55 ± 8.61 ^{ab}
Average daily gain (mg day ⁻¹)	2.78 ± 0.49 ^b	3.24 ± 0.47 ^{ab}	4.43 ± 0.07 ^a	2.51 ± 0.38 ^b	3.61 ± 0.61 ^{ab}
Gastrointestinal weight (mg)	22.08 ± 1.60 ^{ab}	22.28 ± 2.13 ^{ab}	27.03 ± 1.09 ^a	20.30 ± 2.26 ^b	25.17 ± 2.02 ^{ab}
Digestosomatic index (%)	20.64 ± 0.46 ^a	19.77 ± 0.32 ^a	20.40 ± 0.57 ^a	17.92 ± 0.46 ^b	20.53 ± 0.45 ^a

The values in the same row with different superscripts are significantly different ($P < 0.05$).

Table 4

Effects of the experimental diets on digestive enzyme specific activities of total protease (mU mg protein⁻¹), trypsin (μmol *p*-nitroaniline h⁻¹ mg protein⁻¹), chymotrypsin (μmol *p*-nitroaniline h⁻¹ mg protein⁻¹), amylase (μmol maltose h⁻¹ mg protein⁻¹), and lipase (μmol *p*-nitrophenol h⁻¹ mg protein⁻¹) in juvenile Siamese fighting fish at the end of experiment. Data were obtained from triplicate observations.

Parameters	Unmodified	Gamma irradiation	Microwave irradiation	Probiotic	Carbohydrases
Amylase (A)	113.26 ± 2.10 ^b	95.29 ± 7.88 ^{bc}	166.42 ± 1.48 ^a	105.75 ± 10.31 ^b	84.82 ± 1.39 ^c
Total protease	72.33 ± 1.50 ^{ab}	63.30 ± 6.74 ^b	87.11 ± 3.50 ^a	69.09 ± 9.66 ^{ab}	62.03 ± 6.61 ^b
Trypsin (T)	4.20 ± 0.30	3.66 ± 0.07	4.68 ± 0.40	3.91 ± 0.78	3.83 ± 0.34
Chymotrypsin (C)	5.94 ± 0.30	5.98 ± 0.60	6.23 ± 0.39	5.83 ± 0.46	5.96 ± 0.46
Lipase	4.81 ± 0.39	4.69 ± 0.12	3.92 ± 0.02	3.99 ± 0.08	4.39 ± 0.32
T/C ratio	0.71 ± 0.02	0.63 ± 0.07	0.75 ± 0.03	0.63 ± 0.08	0.63 ± 0.04
A/T ratio	27.20 ± 1.64 ^b	26.07 ± 2.28 ^b	36.03 ± 2.87 ^a	20.91 ± 2.82 ^b	24.52 ± 2.29 ^b

The values in the same row with different superscripts are significantly different ($P < 0.05$).

Table 5

Pearson correlation efficiency (r) among different digestive enzyme specific activities and specific growth rate (SGR) of juvenile Siamese fighting fish at the end of experiment. Data were calculated from fish in all treatments ($n = 15$).

	SGR	Amylase	Total protease	Trypsin	Chymotrypsin	Lipase	T/C ratio
SGR	1						
Amylase (A)	0.480	1					
Total protease	0.425	0.851**	1				
Trypsin (T)	0.185	0.635*	0.686**	1			
Chymotrypsin (C)	-0.020	0.253	0.378	0.584*	1		
Lipase	0.259	0.394	0.252	0.467	0.149	1	
T/C ratio	0.272	0.547*	0.466	0.777**	-0.050	0.462	1
A/T ratio	0.535*	0.786**	0.563*	0.028	-0.084	0.142	0.088

* $P < 0.05$, ** $P < 0.01$

Table 6

Effects of the experimental diets on white muscle and body compositions (on wet weight basis) of juvenile Siamese fighting fish at the end of experiment. Data were obtained from triplicate observations.

Parameters	Unmodified	Gamma irradiation	Microwave irradiation	Probiotic	Carbohydrases
White muscle composition					
RNA ($\mu\text{g g}^{-1}$)	4212 \pm 118	4302 \pm 121	4609 \pm 171	4695 \pm 18	4686 \pm 127
Protein (mg g^{-1})	158.21 \pm 3.01 ^a	145.32 \pm 3.59 ^{ab}	150.71 \pm 8.20 ^{ab}	136.68 \pm 7.83 ^b	136.77 \pm 3.25 ^b
Lipid (mg g^{-1})	9.82 \pm 1.01	9.44 \pm 0.35	11.92 \pm 0.67	10.97 \pm 0.89	9.71 \pm 0.90
RNA/protein ratio ($\mu\text{g mg}^{-1}$)	28.75 \pm 1.94	29.62 \pm 0.80	30.53 \pm 0.43	32.81 \pm 0.36	32.17 \pm 2.76
Protein/lipid ratio (mg mg^{-1})	16.44 \pm 1.63	15.46 \pm 0.94	12.66 \pm 0.32	12.53 \pm 0.71	14.38 \pm 1.66
Body composition					
RNA ($\mu\text{g g}^{-1}$)	4349 \pm 243	4269 \pm 120	4565 \pm 240	4445 \pm 215	4357 \pm 317
Protein (mg g^{-1})	151.98 \pm 2.89 ^a	139.60 \pm 3.45 ^{ab}	144.78 \pm 7.88 ^a	131.30 \pm 7.53 ^b	131.39 \pm 3.12 ^b
Lipid (mg g^{-1})	10.14 \pm 0.06	9.56 \pm 1.21	12.28 \pm 1.32	11.43 \pm 0.78	10.63 \pm 1.33
RNA/protein ratio ($\mu\text{g mg}^{-1}$)	28.59 \pm 1.22	30.60 \pm 0.82	31.54 \pm 0.44	33.89 \pm 0.37	33.24 \pm 2.85
Protein/lipid ratio (mg mg^{-1})	15.00 \pm 0.32	14.79 \pm 1.32	12.19 \pm 1.94	11.52 \pm 0.31	12.88 \pm 1.66

The values in the same row with different superscripts are significantly different ($P < 0.05$).